

# THE *Current*

October 22, 2007

Gail Gallessich

## **Scientists Observe Large Galaxy 'Stealing' Gas From Smaller Galaxy**

On Earth, thieves steal everything from diamonds to art to money. In outer space, gas, or fuel for stars, is a commodity worth stealing.

New observations from NASA's Spitzer Space Telescope reveal a distant, massive galaxy in the act of ripping off vast reservoirs of gas – the equivalent of one billion suns – from its smaller, neighbor galaxy. The stolen gas, which has become scorching hot during the heist, will likely cool down and get turned into new stars and planets.

"We may be viewing the larger galaxy in a rare, brief stage of its reincarnation from an old galaxy to a youthful one studded with brilliant stars," said Patrick Ogle of NASA's Spitzer Science Center at the California Institute of Technology in Pasadena. Ogle is the lead author of a new paper on these findings to appear in the Oct. 20 issue of the *Astrophysical Journal*. Robert Antonucci, professor of physics at UC Santa Barbara, is a co-author.

The robber galaxy, called 3C 326 North, is about the size of our Milky Way galaxy, and its victim, 3C 326 South, is about half its mass. They are close enough to perturb each other gravitationally, and might eventually collide. Such galaxy mergers are common in the universe: gas and stars in two nearby galaxies become tangled until they join up into one seamless galaxy. In the case of 3C 326, this is the clearest example yet of large quantities of gas being heated and drained from one

galaxy to another.

"This could be an important phase in galaxy mergers that we are just now witnessing," said Ogle.

Ogle and his colleagues initially set out to study a set of distant galaxies, called radio galaxies, about one billion light-years away. Radio galaxies are named after the radio-emitting jets that scream out of the black holes at their centers. Though these jets are powerful, their black holes are relatively sleepy and don't otherwise give off a lot of energy. When the astronomers scanned 72 galaxies using Spitzer's infrared vision, they noted a handful that were quite unusual.

The most extreme of the bunch, 3C 326 North, was soaking in an enormous amount of hot hydrogen gas, reaching temperatures up to 730 degrees Celsius (1,340 degrees Fahrenheit). This gas, called molecular hydrogen gas because it contains molecules of two hydrogen atoms joined together, is a building block of galaxies, stars and planets. On Earth, it is a potential alternative fuel for cars. Molecular hydrogen gas is invisible to optical telescopes, but when it's heated up it glows at infrared wavelengths that Spitzer can see.

"Hydrogen is by far the predominant element in the universe, yet in its molecular form it has been virtually undetectable until the Spitzer Infrared Space Telescope was launched," said Antonucci of UC Santa Barbara. "The reason is that it is a symmetric molecule, which prevents strong emission or absorption of light at almost any wavelength.

Molecular hydrogen does produce light in a very specific pattern in the infrared part of the spectrum, and suddenly astronomers are finding huge masses of this key gas, providing big surprises in many fields of astronomical study."

Since 3C 326 North is not busy making new stars, it is unusual for it to have this much gas. When the researchers investigated the Spitzer pictures further, they noticed what appeared to be a tail of stars, called a tidal tail, connecting 3C 326 North to 3C 326 South. That was the smoking gun they needed to realize that the pair were interacting, and that 3C 326 North was lifting the gas right off its companion's back.

"The galaxy in question appears to be stripping a large quantity of molecular hydrogen from its neighbor and heating it up," said Ogle. "The supermassive black

hole at the center of the galaxy is digesting a small fraction of the gas and ejecting it in enormous, relativistic jets millions of light-years long."

In addition, the hydrogen gas is hot enough to react with oxygen and form vast quantities of water. This water could potentially be incorporated into planets and comets once the gas has cooled enough to collapse and form new stars and planetary systems.

What will happen to the victim galaxy now that it's bereft of its gas? According to Ogle, the galaxy lost a lot of its fuel, and will no longer produce new stars. However, if the two galaxies do eventually merge, then what belongs to one will belong to the other.

Other authors of this paper include Phil N. Appleton of Caltech and David Whysong of the National Radio Astronomy Observatory, headquartered in Charlottesville, Va.

Related Links

[NASA's Spitzer Space Telescope](#)

---

## **About UC Santa Barbara**

The University of California, Santa Barbara is a leading research institution that also provides a comprehensive liberal arts learning experience. Our academic community of faculty, students, and staff is characterized by a culture of interdisciplinary collaboration that is responsive to the needs of our multicultural and global society. All of this takes place within a living and learning environment like no other, as we draw inspiration from the beauty and resources of our extraordinary location at the edge of the Pacific Ocean.